

CERN-TH/98-177

DFPD-98/TH/25

TUM-HEP-313/98

hep-ph/9806308

June 1998

Next-to-Leading QCD Corrections to $B \rightarrow X_s \gamma$ in Supersymmetry

M. Ciuchini^a, G. Degrossi^b, P. Gambino^c, G.F. Giudice^{d*}

^a *Dipartimento di Fisica, Università di Roma Tre and INFN,
Sezione di Roma III, Via della Vasca Navale, 84, I-00146 Roma, Italy*

^b *Dipartimento di Fisica, Università di Padova, Sezione INFN di Padova,
Via F. Marzolo 8, I-35131 Padua, Italy*

^c *Technische Universität München, Physik Dept., D-85748 Garching, Germany*

^d *Theory Division, CERN CH-1211 Geneva 23, Switzerland*

Abstract

We compute the QCD next-to-leading order matching conditions of the (chromo)-magnetic operators relevant for $B \rightarrow X_s \gamma$ in supersymmetric models with minimal flavour violation. The calculation is performed under the assumption that the charginos and one stop are lighter than all other squarks and the gluino. In the parameter region where a light charged Higgs boson is consistent with measurements of $\text{BR}(B \rightarrow X_s \gamma)$, we find sizeable corrections to the Wilson coefficients. As a consequence, there is a significant reduction of the stop-chargino mass region where the supersymmetric contribution has a large destructive interference with the charged-Higgs boson contribution.

*On leave of absence from INFN, Sez. di Padova, Italy.

1 Introduction

The inclusive decay rate for $B \rightarrow X_s \gamma$ has been first measured by CLEO with the result $\text{BR}(B \rightarrow X_s \gamma) = (2.32 \pm 0.57_{\text{stat}} \pm 0.35_{\text{syst}}) \times 10^{-4}$ [1]. Recently, a preliminary new result based on about 30% more data has been presented by the collaboration, $\text{BR}(B \rightarrow X_s \gamma) = (2.50 \pm 0.47_{\text{stat}} \pm 0.39_{\text{syst}}) \times 10^{-4}$ [2]. The same process has also been measured by ALEPH at LEP, with the result $\text{BR}(B \rightarrow X_s \gamma) = (3.11 \pm 0.80_{\text{stat}} \pm 0.72_{\text{syst}}) \times 10^{-4}$ [3].

There has been significant theoretical effort in refining the prediction of $\text{BR}(B \rightarrow X_s \gamma)$ in the Standard Model (SM). Calculations are now available for the next-to-leading order (NLO) corrections to the anomalous dimensions [4], the matrix elements [5], and the matching conditions of the Wilson coefficients [6, 7], for the leading non-perturbative effects [8], and for the QED corrections [9, 10]. In particular the QCD NLO corrections are important since they reduce the large scale dependence of the leading result [11], which amount to a 30% uncertainty in the leading order (LO) theoretical prediction. Recently, Kagan and Neubert [10] have argued that, at the NLO, the contributions to $\text{BR}(B \rightarrow X_s \gamma)$ from the different Wilson coefficients exhibit a larger scale dependence than the total result, signaling that the theoretical uncertainty may have been underestimated in previous literature [4, 12, 7]. Nevertheless, even considering this effect, they find that the theoretical error can be evaluated to be about 10%. Combining the different theoretical studies, the recent complete analysis in ref. [10] gives the SM prediction $\text{BR}(B \rightarrow X_s \gamma) = (3.29 \pm 0.33) \times 10^{-4} \times \text{BR}(B \rightarrow X_c e \bar{\nu})/10.5\%$.

While the theoretical prediction is made for the total rate of $\text{BR}(B \rightarrow X_s \gamma)$, the experimental data refer only to events with photon energies between 2.2 and 2.7 GeV. The extrapolation from the data to the total rate introduces further theoretical uncertainties from the calculation of the photon energy spectrum [13]. This aspect has been recently emphasized by Kagan and Neubert [10] who have recomputed the spectrum and extrapolated from the CLEO high-energy photon data the total rate $\text{BR}(B \rightarrow X_s \gamma) = (2.66 \pm 0.56_{\text{exp}}^{+0.43}_{-0.48} \text{ th}) \times 10^{-4}$. This is compatible with the SM within one standard deviation.

To summarize, although the SM prediction is still higher than the CLEO result, the discrepancy seems no longer statistically significant. We should however wait for the forthcoming CLEO analysis and future studies at LEP to further clarify the situation. Meanwhile, we use the conservative CLEO upper limit of $\text{BR}(B \rightarrow X_s \gamma) < 4.2 \times 10^{-4}$ at 95% C.L. in our numerical analysis.

The inclusive decay $B \rightarrow X_s \gamma$ is a particularly interesting probe of physics beyond the SM, since it is determined by a flavour-violating loop diagram. Because of this welcome sensitivity on new physics, it is important to determine the predicted rate with sufficient accuracy in a variety of models. In the case of two-Higgs doublet models, results for the next-to-leading order (NLO) QCD corrections have been presented in refs. [7, 14, 15]. In these models, a new contribution arises from charged-Higgs exchange which always increases the SM prediction for $\text{BR}(B \rightarrow X_s \gamma)$. As a consequence, strong limits on the charged-Higgs mass can be derived. These limits are quite dependent on the treatment of theoretical errors and on the amount of discrepancy between theoretical and experimental results [7, 15].

Here we want to extend our previous analysis of QCD NLO effects in $\text{BR}(B \rightarrow X_s \gamma)$ to the case of supersymmetry. Refining the theoretical calculation in supersymmetry [16, 17] (for reviews see ref. [18]) is important because $\text{BR}(B \rightarrow X_s \gamma)$ provides a very stringent constraint on the parameter space of the model. Moreover in supersymmetry it is possible to evade the strong limits on the charged Higgs mass, since the chargino contribution can interfere destructively and reduce $\text{BR}(B \rightarrow X_s \gamma)$ [17]. Our result allows us to give a more reliable estimate of the model parameters necessary to achieve this cancellation.

We have not tried to perform a complete analysis in the most general supersymmetric model, an effort, we feel, that is penalized by the fact that the final result will be very involved and will depend upon many unknown parameters. Instead, we try to focus on what we believe is the most interesting part of the parameter space for $B \rightarrow X_s \gamma$. Specifically, we concentrate on the case in which the flavour violation is completely dictated by the Cabibbo-Kobayashi-Maskawa (CKM) angles (insuring predictability) and in which the charginos and a scalar partner of the top are light (insuring a sizeable new contribution).

The paper is organized as follows. In sect. 2 we define our assumptions on the supersymmetric parameters and discuss the underlying hypotheses. Section 3 contains the analytical results for the supersymmetric corrections to the relevant Wilson coefficients at the matching scale. A numerical analysis of $\text{BR}(B \rightarrow X_s \gamma)$ in supersymmetry at the NLO is presented in sect. 4.

2 Minimal Flavour Violation and Heavy Squark-Gluino Effective Theory

In this section we describe our main theoretical assumptions on the supersymmetric model: *i)* the minimal flavour violation and *ii)* the heavy squark-gluino effective theory that describes chargino, stop, and the SM degrees of freedom with other supersymmetric particles integrated out.

In the supersymmetric extension of the SM with general soft-breaking terms, there are a variety of new sources of flavour violation. The supersymmetry-breaking squark masses and trilinear terms lead to a mismatch in flavour space between quark and squark mass eigenstates. The flavour violation is then described by a large number of mixing angles, which cannot be determined theoretically. All we know is that present measurements and limits on various flavour-violating processes provide quite stringent bounds on these mixing angles (see *e.g.* ref. [19]).

A more predictive case is what we will call *minimal flavour violation*, in which all flavour transitions occur only in the charged-current sector and are determined by the known CKM mixing angles. This is indeed the case in several theoretical schemes in which the communication of the original supersymmetry breaking to the observable particles occurs via flavour-independent interactions. In many of these schemes (especially when supersymmetry breaking occurs at low-energy scales) the departure from the minimal flavour violation hypothesis, caused by quantum effects, is rather small. Our assumption can then be justified in gauge-mediated models [20] (for a review, see ref. [21]) and in certain classes of supergravity theories. At any rate, it plausibly corresponds to the unavoidable flavour violation, present in any supersymmetric model.

Within this class of models, we focus on the special case of interest for $B \rightarrow X_s \gamma$, in which one stop is considerably lighter than the other squarks. In order to preserve the successful fit of the electroweak precision measurements, we assume that the light stop is predominantly right-handed [22]. This assumption can also be theoretically justified by the observation that the renormalization-group evolution of the squark mass parameters indeed pushes the right-handed stop mass to smaller values. Therefore, we concentrate on the case in which the supersymmetry-breaking parameters in the up-squark sector are flavour diagonal (but not necessarily universal) in the basis in which the corresponding up-type quark mass matrix is diagonal. The flavour-violating stop interactions arise only from charged-current effects and

are completely determined by the CKM angles. The stop mass matrix is given by

$$m_t^2 = \begin{pmatrix} m_{\tilde{t}_L}^2 + m_t^2 + (\frac{1}{2} - \frac{2}{3} \sin^2 \theta_W) \cos 2\beta M_Z^2 & m_t(A_t - \mu \cot \beta) \\ m_t(A_t - \mu \cot \beta) & m_{\tilde{t}_R}^2 + m_t^2 + \frac{2}{3} \sin^2 \theta_W \cos 2\beta M_Z^2 \end{pmatrix}. \quad (1)$$

Here $m_{\tilde{t}_R}^2$, $m_{\tilde{t}_L}^2$, and A_t are supersymmetry-breaking parameters, μ is the Higgs mixing mass and $\tan \beta$ is the ratio between the two Higgs vacuum expectation values. We also assume that the charginos are lighter than gluinos and heavy squarks. These approximations considerably simplify the calculation and the final expressions, and they are appropriate to identify the leading supersymmetric contribution.

To implement this scenario we assume the following mass hierarchy

$$\mu_{\tilde{g}} \sim \mathcal{O}(m_{\tilde{g}}, m_{\tilde{q}}, m_{\tilde{t}_1}) \gg \mu_W \sim \mathcal{O}(M_W, m_t, m_{\chi^\pm}, m_{\tilde{t}_2}) \gg m_b \gg \Lambda_{QCD}. \quad (2)$$

Here and in the following μ_W is the mass scale of the charginos ($\chi^\pm$), and of the lighter stop ($\tilde{t}_2$), to be identified with the ordinary electroweak scale. The charged Higgs ($H^\pm$) mass is not constrained. The scale $\mu_{\tilde{g}}$, characteristic of all other strongly-interacting supersymmetric particles (squarks and gluinos), is assumed to be larger, say of the order of the TeV. We compute the QCD NLO corrections to the Wilson coefficients keeping only the first order in an expansion in $\mu_W/\mu_{\tilde{g}}$.

As usual, the presence of different mass scales allows us to use a stack of effective theories, obtained by integrating out of the theory, at each matching scale, the heavy degrees of freedom. The effective theory just below the scale $\mu_{\tilde{g}}$ is particularly simple. The only dimension-five operators involving supersymmetric particles are the chargino (chromo)-magnetic dipole moment and the operator $\bar{t} t \tilde{t}_2^* \tilde{t}_2$, obtained by integrating out the gluino. However, an explicit calculation shows that these operators do not contribute to the NLO matching conditions of magnetic-dipole operators at the scale μ_W . Therefore, to our end, the inclusion of the leading $\mu_W/\mu_{\tilde{g}}$ corrections does not require any new operator other than renormalizable ones.

The next step is the running of the intermediate effective Hamiltonian between $\mu_{\tilde{g}}$ and μ_W . Although the relevant anomalous dimensions are known [23], log resummation is not really needed, given the smallness of the relevant log terms, $\alpha_s(\mu_W)/4\pi \log(\mu_{\tilde{g}}^2/\mu_W^2) \sim 0.01\text{--}0.05$ for typical values of the supersymmetric masses.

Non-resummed NLO effects of heavy supersymmetric particles, including $1/\mu_{\tilde{g}}$ corrections, are simply given by the *one-loop* Feynman diagrams, containing these particles in the loop, which contribute to renormalizable operators in the intermediate effective theory. In other words, we need to compute only corrections involving

gluinos and heavy squarks to the coupling constants appearing in the vertices involving the chargino, $\chi_a^- \bar{b} \tilde{t}_2$, the W vector boson, $W^- \bar{q} q'$, the charged physical, $H^- \bar{q} q'$, and unphysical, $\phi^- \bar{q} q'$, scalars. Heavy $\mathcal{O}(\mu_{\tilde{g}})$ particle effects renormalize the masses of $\mathcal{O}(\mu_w)$ particles, but these effects are reabsorbed through the definition of renormalized masses. We use on-shell masses for squarks. Quark masses are also defined on-shell, except corrections not involving supersymmetric particles, which are subtracted in the $\overline{\text{MS}}$ scheme. This definition simplifies the insertion of supersymmetric contributions into existing RG-evolution formulae, where $\overline{\text{MS}}$ running quark masses are commonly used. Because of our assumption that the light stop is mainly right-handed, the stop mixing angle $\theta_{\tilde{t}}$ is small. For instance, taking the supersymmetry-breaking parameter A_t in eq. (1) of the order of $\mu_{\tilde{g}}$, yields $\theta_{\tilde{t}} \sim \mathcal{O}(\mu_w/\mu_{\tilde{g}})$. In this framework, $1/\mu_{\tilde{g}}$ corrections multiplied by $\sin \theta_{\tilde{t}}$ factors should be regarded as of higher order.

Results for the renormalized vertices in the intermediate effective theory are collected in appendix A. Notice that the $\chi_a^- \bar{b} \tilde{t}_2$ vertex renormalization is infinite. This is not surprising, since supersymmetry and the GIM cancellation are spoiled in the intermediate effective theory, where the heavy squarks and gluinos have been integrated out. This divergence cancels out in the matching of the magnetic operators at the scale μ_w against a corresponding divergence generated by the insertion of the chargino vertex into two-loop diagrams. This cancellation actually provides a check for the results of the two-loop calculation. However, the presence of the divergence calls for a regularization. We choose the naïve dimensional regularization (NDR), to be consistent with the calculation of the anomalous dimension matrix [4]. However, it is known that NDR breaks supersymmetry. In particular, the renormalizations in NDR of the gauge boson and gaugino interactions with matter, as well as the Higgs boson and higgsino interactions, are different and manifestly violate supersymmetry. Supersymmetric Ward identities are restored with appropriate shifts of the gauge and Yukawa couplings in the $\chi_a^- \bar{b} \tilde{t}_2$ vertex [24], denoted as η_Y and η_g in appendix A. These shifts correspond to the difference of using NDR versus dimensional reduction (DR) [25], a regularization that preserves supersymmetry.

The formulae in appendix A are given in terms of few functions of the ratios $m_{\tilde{q}}^2/m_{\tilde{g}}^2$. Notice however that, whenever $m_{\tilde{t}_2}^2/m_{\tilde{g}}^2$ appears, only terms up to $\mathcal{O}(m_{\tilde{t}_2}/m_{\tilde{g}})$ should be retained in the corresponding functions to be consistent with the operator product expansion. Finally, we also keep the leading contribution in the sbottom mixing angle, since the left-right mixing term $(A_b - \mu \tan \beta) m_b$ becomes important for large values of $\tan \beta$.

3 Wilson Coefficients

This section contains the result for the NLO supersymmetric contributions to the Wilson coefficients $C_i^{eff}(\mu_W)$. Following the discussion of the previous section, the calculation of the NLO corrections to $C_7^{eff}(\mu_W)$ and $C_8^{eff}(\mu_W)$ can be divided in two parts: i) the contribution of the heavy particles at the scale $\mu_{\tilde{g}}$ that will appear as renormalization of the coupling constants in the LO diagrams. ii) The contribution of the intermediate-scale particles that requires the computation of the two-loop gluonic corrections to the LO supersymmetric diagrams involving the charginos and the light stop. Concerning the latter two strategies are at hand. One can match matrix elements of operators belonging to a basis obtained enforcing the equations of motion, a procedure that however requires an asymptotic expansion of the relevant diagrams in the external momenta. Alternatively, one can use a larger off-shell operator basis and perform the matching on off-shell matrix elements. In this case, one can use the freedom of the off-shell status to choose a suitable kinematical configuration such that the various Feynman diagrams can be evaluated using ordinary Taylor expansions in the external momenta. This second strategy, already successfully applied by us in ref. [7] to the calculation of the QCD corrections to the matching conditions of the $\Delta B = 1$ magnetic and chromo-magnetic operators in the SM and in two-Higgs doublet models, has also been employed in this calculation. Concerning the technical details we refer to ref. [7].

In the supersymmetric model under consideration we can organize the Wilson coefficients of the operators entering the effective Hamiltonian in the following way:

$$C_i^{eff}(\mu_W) = C_i^{(0)eff}(\mu_W) + \delta^H C_i^{(0)eff}(\mu_W) + \delta^S C_i^{(0)eff}(\mu_W) + \frac{\alpha_s(\mu_W)}{4\pi} \left[C_i^{(1)eff}(\mu_W) + \delta^H C_i^{(1)eff}(\mu_W) + \delta^S C_i^{(1)eff}(\mu_W) \right] \quad (3)$$

where $C_i^{(k)eff}(\mu_W)$ represents the SM contribution ($k = 0, 1$), $\delta^H C_i^{(k)eff}(\mu_W)$ the additional terms present in a two-Higgs doublet model, while $\delta^S C_i^{(k)eff}(\mu_W)$ contains the contribution from supersymmetric particles. Explicit expressions for $C_i^{(k)eff}(\mu_W)$ and $\delta^H C_i^{(k)eff}(\mu_W)$ can be found in sections 4 and 5 of ref. [7]¹. Concerning the remaining contributions, the LO $\delta^S C_i^{(0)eff}(\mu_W)$ terms are given by [16]

$$\begin{aligned} \delta^S C_i^{(0)eff}(\mu_W) &= 0 \quad i = 1, \dots, 6 \\ \delta^S C_{7,8}^{(0)eff}(\mu_W) &= \sum_{j=1,2} \left[\frac{2}{3} \frac{M_W^2}{\tilde{m}^2} \tilde{V}_{j1}^2 F_{7,8}^{(1)}(z_j) + \frac{\tilde{U}_{j2}}{\sqrt{2} \cos \beta} \frac{M_W}{m_{\chi_j}} \tilde{V}_{j1} F_{7,8}^{(3)}(z_j) \right] \end{aligned}$$

¹In ref. [7] $\delta^H C_i^{(k)eff}$ are indicated as $\delta C_i^{(k)eff}$.

$$\begin{aligned} & -\frac{2}{3}t_{1j}^2 \frac{M_W^2}{m_{\tilde{t}_1}^2} F_{7,8}^{(1)}(y_{1j}) - \frac{\tilde{U}_{j2}}{\sqrt{2} \cos \beta} \frac{M_W}{m_{\chi_j}} t_{1j} \cos \theta_{\tilde{t}} F_{7,8}^{(3)}(y_{1j}) \\ & -\frac{2}{3}t_{2j}^2 \frac{M_W^2}{m_{\tilde{t}_2}^2} F_{7,8}^{(1)}(y_{2j}) - \frac{\tilde{U}_{j2}}{\sqrt{2} \cos \beta} \frac{M_W}{m_{\chi_j}} t_{2j} \sin \theta_{\tilde{t}} F_{7,8}^{(3)}(y_{2j}) \end{aligned} \quad (4)$$

where the functions $F_{7,8}^{(1)}$ are defined in eqs. (29)–(30) of ref. [7] and

$$\begin{aligned} F_7^{(3)}(x) &= \frac{5-7x}{6(x-1)^2} + \frac{x(3x-2)}{3(x-1)^3} \ln x \\ F_8^{(3)}(x) &= \frac{1+x}{2(x-1)^2} - \frac{x}{(x-1)^3} \ln x. \end{aligned} \quad (5)$$

We have defined

$$t_{1j} = \tilde{V}_{j1} \cos \theta_{\tilde{t}} - \tan \theta_{\tilde{t}} Y_j; \quad t_{2j} = \tilde{V}_{j1} \sin \theta_{\tilde{t}} + Y_j; \quad (6)$$

$$z_j = \frac{\tilde{m}^2}{m_{\chi_j}^2}; \quad y_{kj} = \frac{m_{\tilde{t}_k}^2}{m_{\chi_j}^2}; \quad Y_j = \frac{\tilde{V}_{j2} \cos \theta_{\tilde{t}}}{\sqrt{2} \sin \beta} \frac{\bar{m}_t(\mu_w)}{M_W}, \quad (7)$$

where $\bar{m}_t(\mu_w)$ is the top-quark running mass at the scale μ_w .

The stop eigenstates $\tilde{t}_1 = \cos \theta_{\tilde{t}} \tilde{t}_L + \sin \theta_{\tilde{t}} \tilde{t}_R$ and $\tilde{t}_2 = -\sin \theta_{\tilde{t}} \tilde{t}_L + \cos \theta_{\tilde{t}} \tilde{t}_R$ have mass eigenvalues $m_{\tilde{t}_1}$ and $m_{\tilde{t}_2}$ and we have taken all other squarks to be degenerate with mass $\tilde{m}$. The two matrices $\tilde{U}$ and $\tilde{V}$ diagonalize the chargino mass matrix according to (M is the weak gaugino mass)

$$\tilde{U} \begin{pmatrix} M & M_W \sqrt{2} \sin \beta \\ M_W \sqrt{2} \cos \beta & \mu \end{pmatrix} \tilde{V}^{-1} \quad (8)$$

and are assumed to be real. Notice that in eq. (4) and henceforth the scalar quark masses are understood as on-shell.

The last ingredients needed for a complete NLO calculation in our supersymmetric model are $\delta^S C_i^{(1)eff}(\mu_w)$. For the current-current and penguins operators we have

$$\delta^S C_i^{(1)eff}(\mu_w) = 0 \quad i = 1, 2, 3, 5, 6 \quad (9)$$

$$\delta^S C_4^{(1)eff}(\mu_w) = \frac{M_W^2}{m_{\tilde{t}_2}^2} \sum_{j=1,2} t_{2j}^2 E_\chi(y_{2j}) \quad (10)$$

with [26]

$$E_\chi(x) = \frac{x(11-7x+2x^2)}{18(x-1)^3} - \frac{x}{3(x-1)^4} \ln x. \quad (11)$$

The $O(\alpha_s)$ corrections to the coefficient of the (chromo-)magnetic operator can be divided into 4 pieces

$$\begin{aligned}\delta^S C_{7,8}^{(1)eff}(\mu_W) &= \delta^X C_{7,8}^{(1)eff}(\mu_W) + \delta^W C_{7,8}^{(1)eff}(\mu_W) + \delta^{\phi_1} C_{7,8}^{(1)eff}(\mu_W) \\ &+ \delta^{\phi_2} C_{7,8}^{(1)eff}(\mu_W) .\end{aligned}\quad (12)$$

Here $\delta^X C_{7,8}^{(1)eff}$ represents the chargino contribution while $(\delta^W, \delta^{\phi_1}, \delta^{\phi_2}) C_{7,8}^{(1)eff}$ take into account the renormalization effects due to the $O(\mu_{\tilde{g}})$ heavy particles in the W , physical and unphysical charged scalar couplings, respectively. We find

$$\begin{aligned}\delta^X C_7^{(1)eff}(\mu_W) &= \sum_{j=1,2} \left[t_{2j}^2 \frac{M_W^2}{m_{\tilde{t}_2}^2} \left(G_7^{\chi,1}(y_{2j}) + \Delta_7^{\chi,1}(y_{2j}) \ln \frac{\mu_W^2}{m_{\chi_j}^2} - \frac{4}{9} E_\chi(y_{2j}) \right) \right. \\ &+ \frac{\tilde{U}_{j2} \sin \theta_{\tilde{t}}}{\sqrt{2} \cos \beta} \frac{M_W}{m_{\chi_j}} \left(t_{2j} G_7^{\chi,2}(y_{2j}) + t_{2j} \Delta_7^{\chi,2}(y_{2j}) \ln \frac{\mu_W^2}{m_{\chi_j}^2} \right. \\ &\left. \left. - \frac{4}{3} Y_j R_s F_7^{(3)}(y_{2j}) \right) - \frac{8}{9} Y_j \frac{M_W^2}{m_{\tilde{t}_2}^2} t_{2j} (R_s + R_b) F_7^{(1)}(y_{2j}) \right] \quad (13)\end{aligned}$$

$$\delta^X C_8^{(1)eff}(\mu_W) = \delta^X C_7^{(1)eff}(\mu_W) \left(7 \rightarrow 8, -\frac{4}{9} E_\chi(y_{2j}) \rightarrow -\frac{1}{6} E_\chi(y_{2j}) \right). \quad (14)$$

Eq. (14) means that $\delta^X C_8^{(1)eff}(\mu_W)$ can be obtained from the r.h.s. of eq. (13) by replacing the index 7 with the index 8 in the various functions and by substituting the term $-4/9 E_\chi(y_{2j})$ with $-1/6 E_\chi(y_{2j})$. In eqs. (13)–(14) we have used

$$\begin{aligned}G_7^{\chi,1}(x) &= -\frac{8}{9} F_7^{(1)}(x) \left(\Delta_{t,s}^{(1)} + \Delta_{t,b}^{(1)} - 1 + 3 \ln \frac{m_{\tilde{g}}^2}{m_{\chi_j}^2} \right) \\ &+ \frac{x(85 - 347x + 526x^2)}{243(1-x)^3} + \frac{4x^2(-8 + 13x + 6x^2)}{9(x-1)^4} \text{Li}_2 \left(1 - \frac{1}{x} \right) \\ &- \frac{4x(20 - 126x + 144x^2 + 39x^3)}{81(x-1)^4} \ln x + \frac{2x^2(21x - 10)}{9(x-1)^4} \ln^2 x \quad (15)\end{aligned}$$

$$\begin{aligned}G_7^{\chi,2}(x) &= -\frac{4}{3} F_7^{(3)}(x) \left(\Delta_{t,s}^{(1)} + \Delta_b^{(1)} + \Delta_b^{(2)} - 2 \right) - \frac{16(3-7x)x}{9(x-1)^3} \text{Li}_2 \left(1 - \frac{1}{x} \right) \\ &+ \frac{4(3x-5)}{9(x-1)^2} - \frac{4(4-30x+40x^2)}{9(x-1)^3} \ln x - \frac{16(1-3x)x}{9(x-1)^3} \ln^2 x \quad (16)\end{aligned}$$

$$G_8^{\chi,1}(x) = -\frac{8}{9} F_8^{(1)}(x) \left(\Delta_{t,s}^{(1)} + \Delta_{t,b}^{(1)} - 1 + 3 \ln \frac{m_{\tilde{g}}^2}{m_{\chi_j}^2} \right)$$

$$\begin{aligned}
& - \frac{x(1210 - 437x - 1427x^2)}{648(x-1)^3} - \frac{x^2(49 + 46x + 9x^2)}{12(x-1)^4} \text{Li}_2\left(1 - \frac{1}{x}\right) \\
& - \frac{x(85 - 603x - 387x^2 + 78x^3)}{108(x-1)^4} \ln x - \frac{13x^2}{3(x-1)^4} \ln^2 x
\end{aligned} \tag{17}$$

$$\begin{aligned}
G_8^{\chi,2}(x) &= -\frac{4}{3}F_8^{(3)}(x) \left(\Delta_{t,s}^{(1)} + \Delta_b^{(1)} + \Delta_b^{(2)} - 2 \right) - \frac{4x(3+4x)}{3(x-1)^3} \text{Li}_2\left(1 - \frac{1}{x}\right) \\
& - \frac{61 - 39x}{12(x-1)^2} - \frac{(7 - 60x - 14x^2)}{6(x-1)^3} \ln x - \frac{14x}{3(x-1)^3} \ln^2 x
\end{aligned} \tag{18}$$

$$R_i = 3 \ln \frac{\mu_W^2}{m_{\tilde{g}}^2} + \Delta_{t,i}^{(2)} - 1 \tag{19}$$

$$\Delta_7^{\chi,1}(x) = \frac{32}{27} \left(F_8^{(1)}(x) - 3F_7^{(1)}(x) \right); \quad \Delta_7^{\chi,2}(x) = \frac{16}{9} \left(F_8^{(3)}(x) - 3F_7^{(3)}(x) \right) \tag{20}$$

$$\Delta_8^{\chi,1}(x) = -\frac{28}{9}F_8^{(1)}(x); \quad \Delta_8^{\chi,2}(x) = -\frac{14}{3}F_8^{(3)}(x) \tag{21}$$

The various functions Δ appearing in eqs. (15)–(19) contain the effect of the renormalization of the chargino-stop-quark vertex due to the $\mathcal{O}(\mu_{\tilde{g}})$ heavy particles. Their explicit expressions are given in appendix A. In the same equations the terms not proportional to the one-loop functions $F_{7,8}^{(1,3)}$ represent the μ -independent part of the contributions coming from two-loop squark-chargino diagrams. The μ_W dependence in eq. (13) and eq. (14) satisfies the relation

$$\begin{aligned}
\frac{1}{2}\gamma_0^m \bar{m}_t \frac{\partial C_i^{(0)}}{\partial \bar{m}_t} &+ \frac{1}{2} \sum_{l=1}^8 \gamma_{li}^{(0)eff} C_l^{(0)} = \\
& \sum_{j=1}^2 \left\{ \frac{M_W^2}{m_{\tilde{t}_2}^2} \left(t_{2j}^2 \Delta_i^{\chi,1}(y_{2j}) - \frac{16}{3} t_{2j} Y_j F_i^{(1)}(y_{2j}) \right) \right. \\
& \left. + \frac{\tilde{U}_{j2} \sin \theta_{\tilde{t}} M_W}{\sqrt{2} \cos \beta m_{\chi_j}} \left(t_{2j} \Delta_i^{\chi,2}(y_{2j}) - 4 Y_j F_i^{(3)}(y_{2j}) \right) \right\}
\end{aligned} \tag{22}$$

that ensures that physical observables are independent of μ_W to $\mathcal{O}(\alpha_s)$. In eq. (22) $\gamma_0^m = 8$ is the LO anomalous dimension of the top mass.

Finally, we report the effects of $\mathcal{O}(\mu_{\tilde{g}})$ heavy particles in diagrams involving W , physical and unphysical charged scalars exchanges. As previously discussed, in our approximation they are introduced as a renormalization of the relevant couplings

and the corresponding contributions are given by

$$\delta^W C_7^{(1)eff}(\mu_W) = \frac{4}{3} (W_t^b + W_t^s) G_7^W(tw) - \frac{23}{27} (W_c^b + W_c^s) \quad (23)$$

$$\delta^W C_8^{(1)eff}(\mu_W) = \frac{4}{3} (W_t^b + W_t^s) G_8^W(tw) - \frac{4}{9} (W_c^b + W_c^s) \quad (24)$$

$$\begin{aligned} \delta^{\phi_1} C_{7,8}^{(1)eff}(\mu_W) &= \frac{4}{9 \tan^2 \beta} (H_t^s + H_t^b) F_{7,8}^{(1)}(th) \\ &+ \frac{4}{3} (H_t^s + H_b) F_{7,8}^{(2)}(th) \end{aligned} \quad (25)$$

$$\delta^{\phi_2} C_{7,8}^{(1)eff}(\mu_W) = \frac{4}{9} (U_t^s + U_t^b) F_{7,8}^{(1)}(tw) - \frac{4}{3} (U_t^s + U_b) F_{7,8}^{(2)}(tw) \quad . \quad (26)$$

Here $tw = \bar{m}_t(\mu_W)^2/M_W^2$, $th = \bar{m}_t(\mu_W)^2/M_{H^\pm}^2$,

$$\begin{aligned} G_7^W(x) &= -\frac{23 - 67x + 50x^2}{36(x-1)^3} + \frac{x(2 - 7x + 6x^2)}{6(x-1)^4} \ln x; \\ G_8^W(x) &= -\frac{4 - 5x - 5x^2}{12(x-1)^3} + \frac{x(1 - 2x)}{2(x-1)^4} \ln x, \end{aligned} \quad (27)$$

the functions $F_{7,8}^{(2)}$ are defined in eqs. (54)–(55) of ref. [7], and the renormalization functions W, H, U are given in appendix A.

4 Numerical Analysis

In this section we present a numerical analysis of $\text{BR}(B \rightarrow X_s \gamma)$, with special attention to the comparison between LO and NLO results in the supersymmetric model. We focus on the situation in which our approximation is adequate, *i.e.* light stop and charginos, and show how the strong bounds on the charged Higgs mass can be relaxed.

We use the same numerical inputs and conventions as in ref. [7], except for the photon-energy cut off $E_\gamma > (1 - \delta)m_b/2$, which is now chosen to be $\delta = 0.90$ (see discussion in ref. [10]). In our numerical analysis we also include the leading logarithmic electromagnetic effects as computed in refs. [9, 10]. They amount to a decrease of the calculated branching ratio of about 7% to 8%. We neglect all other two-loop electroweak effects which appear to be very small [27].

Light supersymmetric particles can affect the measured values of the $B^0 - \bar{B}^0$ mixing ΔM_B and of the CP-violating ϵ_K parameter and, ultimately, the extraction of the CKM angles, which enter the calculation of $\text{BR}(B \rightarrow X_s \gamma)$. Within our

assumptions, it is not difficult to compute this effect. The dominant supersymmetric contributions to ΔM_B and ϵ_K come from box diagrams with chargino/light stop and charged Higgs/top quark exchange. Under the assumption of minimal flavour violation, these contributions have the same structure of CKM angles as the top-quark box diagram in the SM. Since the dominant effects on ΔM_B and ϵ_K in the SM indeed come from the top quark, the CKM-angle dependence is the same for all contributions, and the effect of supersymmetry is parametrized by

$$R = 1 + \frac{\Delta_{SUSY}}{\Delta_{SM}}. \quad (28)$$

Here Δ_{SM} is the amplitude of the top-quark box diagram, and Δ_{SUSY} is the amplitude of the chargino/light stop and charged Higgs/top quark box diagrams, whose analytical expressions are given in the appendix B. The crucial point is that R is independent of CKM angles. In the whole supersymmetric parameter space, R turns out to be larger than one.

In terms of the Wolfenstein parameters η and ρ , the CKM structure of ΔM_B and ϵ_K in supersymmetry is

$$\Delta M_B \sim |V_{td}^* V_{tb}|^2 R \sim [(1 - \rho)^2 + \eta^2] R \quad (29)$$

$$\epsilon_K \sim |\text{Im}(V_{td}^* V_{ts})^2| R \sim \eta(1 - \rho)R. \quad (30)$$

The new physics contribution can be effectively reabsorbed in the Wolfenstein parameters $(1 - \rho)\sqrt{R}$ and $\eta\sqrt{R}$. This means that the “true” values of η and ρ are related to the values η_{SM} and ρ_{SM} extracted from the usual SM fit by

$$\eta = \frac{\eta_{SM}}{\sqrt{R}} \quad (31)$$

$$\rho = 1 - \frac{(1 - \rho_{SM})}{\sqrt{R}}. \quad (32)$$

We have checked that eqs. (31)–(32) give an extremely good approximation of a complete numerical fit. The combination of the CKM angles entering the evaluation of $\text{BR}(B \rightarrow X_s \gamma)$ is

$$X \equiv \left| \frac{V_{ts}^* V_{tb}}{V_{cb}} \right|^2 = 1 + (2\rho - 1)\lambda^2 = 1 + \left[1 - \frac{2}{\sqrt{R}}(1 - \rho_{SM}) \right] \lambda^2, \quad (33)$$

where $\lambda = |V_{us}|$ does not depend on the supersymmetric masses. Therefore, in the presence of supersymmetry, the value of X is related to the usual SM input X_{SM} by

$$X = X_{SM} + (\lambda^2 + 1 - X_{SM}) \left(1 - \frac{1}{\sqrt{R}} \right). \quad (34)$$

This is the value of X to be used in a supersymmetric analysis of $\text{BR}(B \rightarrow X_s \gamma)$.

It is known that the chargino/stop contribution to the Wilson coefficient C_7 can interfere with the SM and charged Higgs contributions either constructively or destructively, depending on the parameters choice. The case of destructive interference is particularly interesting, since it is possible to make a light charged Higgs consistent with the measurement of $\text{BR}(B \rightarrow X_s \gamma)$. For illustration, we choose the supersymmetric parameters corresponding to a strong cancellation between the chargino and charged-Higgs contributions, and plot in fig. 1 the predicted $\text{BR}(B \rightarrow X_s \gamma)$ as a function of $\mu_{\tilde{g}}$. The LO result in supersymmetry is compared with the result obtained in the limit $\mu_{\tilde{g}} \rightarrow \infty$ (denoted by **LOdecoupled** in fig. 1). This shows how the asymptotic behaviour is reached, as $\mu_{\tilde{g}}$ grows. The NLO result is also shown in fig. 1. In this case, we cannot define an asymptotic behaviour, as the NLO branching ratio hardly decreases with $\mu_{\tilde{g}}$ and actually starts rising again for $\mu_{\tilde{g}} > 0.7$ TeV. This non-decoupling effect is not surprising [28]. It appears because, at the NLO, the chargino vertex contains terms proportional to $\log(\mu_{\tilde{g}}/\mu_w)$. Indeed, the chargino coupling constant is related to the ordinary gauge and Yukawa couplings only by supersymmetry. In the effective theory below $\mu_{\tilde{g}}$, supersymmetry is explicitly broken and the renormalization of the chargino vertex develops a logarithm of the large supersymmetry-breaking scale. Notice the important effect of the NLO corrections calculated here. For the supersymmetric parameters of fig. 1 and for $\mu_{\tilde{g}} = 1$ TeV, the LO branching ratio is 1.9×10^{-4} , while the NLO result is 3.7×10^{-4} . Had we neglected in the NLO analysis our new supersymmetric contributions to the matching conditions, the result would have been 2.4×10^{-4} . The large shift induced by the NLO corrections to $C_{7,8}^{eff}(\mu_w)$ is mainly due by the large non-decoupling logarithms of $\mu_{\tilde{g}}/\mu_w$. Finally, in fig. 1 we also show the LO and NLO results obtained in a two-Higgs doublet model with charged-Higgs mass equal to the supersymmetric case. The light charged Higgs (taken with a mass of 100 GeV in fig. 1) is clearly incompatible with the measurement of $\text{BR}(B \rightarrow X_s \gamma)$, in the absence of an appropriate chargino contribution.

In order to assess the impact of the different contributions, it is also interesting to consider the NLO Wilson coefficients $C_{7,8}^{eff}$ at the weak scale $\mu_w = M_W$ in the supersymmetric configuration chosen in fig. 1 with all heavy masses set to 1 TeV, and compare them with the corresponding quantities in the SM and the two-Higgs doublet model. The results are shown in the table 1. Notice that, in the supersymmetric case, the QCD corrections to C_7^{eff} amount to about 60%. Such a large correction is clearly related to the fact that the LO approximate cancellation among

different contributions is partially spoiled at the NLO level. Indeed, in the case under consideration, the NLO effects increase the SM contribution to $|C_7^{eff}(M_W)|$ by almost 10% and decrease the charged-Higgs and the chargino contributions by about 20% and 30%, respectively. This situation of partial cancellation and enhanced sensitivity to NLO correction is actually the most interesting phenomenologically. It is under this condition that a light charged-Higgs mass is still allowed and that our calculation is essential.

	$C_7^{eff}(M_W)$	$C_8^{eff}(M_W)$
LO SM	-0.198	-0.098
NLO SM	-0.220	-0.119
LO 2HDM	-0.529	-0.336
NLO 2HDM	-0.493	-0.326
LO SUSY	-0.143	-0.141
NLO SUSY	-0.229	-0.183

Table 1: Wilson coefficients evaluated at the scale M_W in the Standard Model (SM), the two-Higgs doublet model (2HDM) and supersymmetry (SUSY) at LO and at NLO for $\tan\beta = 1$, $M_{H^\pm} = m_{\tilde{t}_2} = m_{\chi_2} = 100$ GeV, $m_{\chi_1} = 300$ GeV, $\theta_{\tilde{t}} = -\pi/10$, $A_b = A_t$; all other squarks and the gluino are degenerate with mass of 1 TeV. The lighter chargino is predominantly higgsino.

We now want to investigate the region of parameter space in which we can significantly relax the bound on the charged Higgs mass from $\text{BR}(B \rightarrow X_s \gamma)$. We have scanned over the relevant supersymmetric parameters, assuming a stop mixing angle less than $\pi/10$ in absolute value, consistently with our assumption of a mainly right-handed light stop. In fig. 2 we show, for $\tan\beta = 2$ and 4, the maximum values of the lighter chargino and lighter stop masses for which a charged Higgs of 100 GeV is consistent with the 95% C.L. CLEO limit of 4.2×10^{-4} . We have chosen this value to give a conservative estimate. With improved experimental results and revised analyses of the photon energy spectrum, this bound may become significantly more restrictive. The effect of the NLO corrections is to make the upper bounds on the chargino and stop masses quite more stringent. In the same figure we also present the results obtained using the renormalization group evolution and the SM and charged-Higgs contributions to the Wilson coefficients at the NLO, while the

purely supersymmetric effects on $C_{7,8}^{eff}(\mu_w)$ are evaluated at the LO. This is to show the impact of the calculation presented here. As seen from the figure, the effect of the QCD corrections to the supersymmetric contribution to $C_{7,8}^{eff}(\mu_w)$ is very significant.

5 Conclusions

In this paper we have computed the QCD NLO corrections to the matching conditions of the Wilson coefficients relevant for $\text{BR}(B \rightarrow X_s \gamma)$. We have used some theoretical assumptions to simplify the result and to concentrate on the most relevant part of the supersymmetric parameter space. We have assumed minimal flavour violation or, in other words, that the flavour-violating interactions in supersymmetry are primarily in the charged-current sector and are determined by the CKM angles. This is often a good approximation in a variety of models. At worst, our result represents an unavoidable contribution to be added to new effects coming from other sources of flavour violation. This assumption allow us also to limit the number of unknown parameters.

We have focused on the case in which the purely supersymmetric contribution is sizeable and can compensate the effect of a light charged-Higgs boson. For this reason, we have assumed that charginos and a mainly right-handed stop are considerably lighter than the other squarks and the gluino. This assumption has allowed us to use an effective theory, in which the heavy particles are integrated out, retaining only the first term in an expansion in the ratio between light and heavy masses.

In sect. 3 we have presented the analytic results for the NLO matching conditions of the Wilson coefficients in supersymmetry, under the approximations stated above. We have also performed a numerical investigation of the supersymmetric parameter region in which there is a sizeable cancellation among various different terms. In this region the impact of the NLO corrections to the supersymmetric contribution to the Wilson coefficients is very significant because of two conspiring effects. On one side there is a large renormalization of the one-loop supersymmetric contribution, mostly coming from logarithms of the ratio between the high and the intermediate mass scales. On the other side, the effect of the large NLO corrections to the supersymmetric contribution is greatly enhanced whenever there is an approximate cancellation at the LO.

We would like to thank A. Buras, G. Martinelli, M. Misiak, M. Neubert, and L. Silvestrini for useful discussions. Right before submitting this paper, we learned from M. Misiak and J. Urban that they have reproduced our results in the limit of infinite gluino mass.

Appendix A

In this appendix we present the results for the renormalized vertices in the effective theory at scales between $\mu_{\tilde{g}}$ and μ_w . In deriving the various corrections we exploited the fact that, in our framework, the light stop is mainly right-handed and therefore we can neglect terms $O(\sin \theta_{\tilde{t}} \mu_w / \mu_{\tilde{g}})$. However in the formulae below we do not expand functions of $m_{\tilde{t}_2}^2 / m_{\tilde{g}}^2$ and we keep the explicit $\sin^2 \theta_{\tilde{t}}$ terms, because we feel it will be easier for the reader to understand the origin of the different terms.

W

The renormalized $\bar{d} u W$ vertex is

$$(-i) \frac{g}{\sqrt{2}} V_{ud} \gamma^\mu a_- \left(1 + \frac{\alpha_s}{3\pi} W_u^d \right)$$

where

$$W_u^d = \frac{1}{2} \left(\cos^2 \theta_{\tilde{u}} W[x_1, w_1] + \sin^2 \theta_{\tilde{u}} W[x_1, w_2] \right)$$

with $x_1 = m_{\tilde{d}_1}^2 / m_{\tilde{g}}^2$, $w_i = m_{\tilde{u}_i}^2 / m_{\tilde{g}}^2$ and

$$W[x, y] = \frac{x + y - 2xy}{(x-1)(y-1)} + \frac{x^3 - 2xy + x^2y}{(x-1)^2(x-y)} \ln x + \frac{2xy - xy^2 - y^3}{(y-1)^2(x-y)} \ln y.$$

For equal masses $W[x, x] = 0$.

Unphysical scalar

The renormalized $\bar{d} t \phi^+$ vertex is

$$(-i) \frac{g}{\sqrt{2} M_w} V_{td} \left[m_d a_- \left(1 + \frac{\alpha_s}{3\pi} U_d \right) - m_t a_+ \left(1 + \frac{\alpha_s}{3\pi} U_t^d \right) \right]$$

where

$$U_d = \frac{1}{2} \left(H_1[x_1] - \cos^2 \theta_{\tilde{t}} H_1[u_1] - \sin^2 \theta_{\tilde{t}} H_1[u_2] \right)$$

$$\begin{aligned}
& -2 \frac{A_d - \mu \tan \beta}{m_{\tilde{g}}} H_2[x_1, x_2] \\
& + 2 \frac{A_d - \mu \tan \beta}{m_{\tilde{g}}} \left(\cos^2 \theta_{\tilde{t}} H_2[u_1, x_2] + \sin^2 \theta_{\tilde{t}} H_2[u_2, x_2] \right), \\
U_t^d = & -\frac{1}{2} \left(H_1[x_1] - \cos^2 \theta_{\tilde{t}} H_1[u_1] - \sin^2 \theta_{\tilde{t}} H_1[u_2] \right) \\
& - 2 \frac{A_u - \mu \cot \beta}{m_{\tilde{g}}} H_2[u_1, u_2] \\
& + 2 \frac{A_u - \mu \cot \beta}{m_{\tilde{g}}} \left(\cos^2 \theta_{\tilde{t}} H_2[u_2, x_1] + \sin^2 \theta_{\tilde{t}} H_2[u_1, x_1] \right).
\end{aligned}$$

with $x_2 = m_{\tilde{d}_2}^2/m_{\tilde{g}}^2$, $u_i = m_{\tilde{t}_i}^2/m_{\tilde{g}}^2$ and

$$\begin{aligned}
H_1[x] &= \frac{1}{1-x} + \frac{2x-x^2}{(1-x)^2} \ln x \\
H_2[x, y] &= \frac{x}{(1-x)(x-y)} \ln x + \frac{y}{(1-y)(y-x)} \ln y
\end{aligned}$$

Physical scalar

We write the renormalized vertex $\bar{d} t h^+$ as

$$\frac{ig}{\sqrt{2} M_W} V_{td} \left[\tan \beta m_d a_- \left(1 + \frac{\alpha_s}{3\pi} H_d \right) + \cot \beta m_t a_+ \left(1 + \frac{\alpha_s}{3\pi} H_t^d \right) \right]$$

where

$$\begin{aligned}
H_d &= \frac{1}{2} \left(H_1[x_1] - \cos^2 \theta_{\tilde{t}} H_1[u_1] - \sin^2 \theta_{\tilde{t}} H_1[u_2] \right) \\
& - 2 \frac{A_d - \mu \tan \beta}{m_{\tilde{g}}} H_2[x_1, x_2] \\
& + 2 \frac{A_d + \mu \cot \beta}{m_{\tilde{g}}} \left(\cos^2 \theta_{\tilde{t}} H_2[u_1, x_2] + \sin^2 \theta_{\tilde{t}} H_2[u_2, x_2] \right), \\
H_t^d &= -\frac{1}{2} \left(H_1[x_1] - \cos^2 \theta_{\tilde{t}} H_1[u_1] - \sin^2 \theta_{\tilde{t}} H_1[u_2] \right) \\
& - 2 \frac{A_u - \mu \cot \beta}{m_{\tilde{g}}} H_2[u_1, u_2] \\
& + 2 \frac{A_u + \mu \tan \beta}{m_{\tilde{g}}} \left(\cos^2 \theta_{\tilde{t}} H_2[u_2, x_1] + \sin^2 \theta_{\tilde{t}} H_2[u_1, x_1] \right).
\end{aligned}$$

Chargino

Unlike the case of the W and of physical and unphysical scalars, in the chargino sector the $O(\alpha_s)$ corrections due to gluon and gluino exchanges are not separately finite. Therefore, as shown below, the renormalization of the chargino vertex due to the $O(\mu_{\tilde{g}})$ heavy particles will contain $1/\epsilon$ poles ($\epsilon = (4 - n)/2$, with n the dimension of the space-time). The cancellation of these poles against similar terms coming from two-loop diagrams in which a gluon is present provides a check of our calculation. We write the corrected $\bar{\chi}_a d \tilde{t}_2$ vertex as

$$(-ig) V_{td} \left\{ -\sin \theta_{\tilde{t}} \tilde{U}_{a2} \frac{m_d}{\sqrt{2} M_W \cos \beta} a_+ \left(1 + \frac{\alpha_s}{3\pi} C_d \right) + \left[\cos \theta_{\tilde{t}} \tilde{V}_{a2} \frac{m_t}{\sqrt{2} M_W \sin \beta} a_- \left(1 + \frac{\alpha_s}{3\pi} C_{t,d}^a \right) + \sin \theta_{\tilde{t}} \tilde{V}_{a1} a_- \left(1 + \frac{\alpha_s}{3\pi} C_{t,d}^b \right) \right] \right\} \quad (\text{A1})$$

with ($\bar{\mu}$ is the 't-Hooft mass)

$$\begin{aligned} C_d &= \frac{3}{2\epsilon} - \frac{3}{2} \ln \left(\frac{m_{\tilde{g}}^2}{\bar{\mu}^2} \right) + \Delta_d^{(1)} + \Delta_d^{(2)} + \eta_Y \\ C_{t,d}^a &= \frac{3}{2\epsilon} - \frac{3}{2} \ln \left(\frac{m_{\tilde{g}}^2}{\bar{\mu}^2} \right) + \Delta_{t,d}^{(1)} + \Delta_{t,d}^{(2)} + \eta_Y \\ C_{t,d}^b &= -\frac{3}{2\epsilon} + \frac{3}{2} \ln \left(\frac{m_{\tilde{g}}^2}{\bar{\mu}^2} \right) + \Delta_{t,d}^{(1)} + \eta_g. \end{aligned} \quad (\text{A2})$$

The $\Delta_{t(d)}^{(i)}$ functions in eq. (A2), also appearing in eqs. (15)–(19), are given by

$$\begin{aligned} \Delta_d^{(1)} &= -\frac{3}{4} - \frac{1}{2} H_1[x_2] \\ \Delta_{t,d}^{(1)} &= -\frac{3}{4} - \frac{1}{2} H_1[x_1] \\ \Delta_d^{(2)} &= \frac{5}{2} + \left(1 - \cot \theta_{\tilde{t}} \frac{m_t m_{\tilde{g}}}{m_{d_2}^2} \right) H_3[x_2] + \frac{1}{2} H_1[x_1] + \frac{1}{2} H_1[x_2] \\ &\quad - 2 \frac{A_d - \mu \tan \beta}{m_{\tilde{g}}} H_2[x_1, x_2] \\ \Delta_{t,d}^{(2)} &= \frac{5}{2} + H_3[x_1] + \frac{1}{2} H_1[u_1] + \frac{1}{2} H_1[u_2] - 2 \frac{A_u - \mu \cot \beta}{m_{\tilde{g}}} H_2[u_1, u_2] \end{aligned}$$

where

$$H_3[x] = \frac{2x}{1-x} \ln x.$$

In eq. (A2) the factors $\eta_Y = -3/2$ and $\eta_g = -1/2$ are induced by the fact that the $\overline{\text{MS}}$ renormalization does not preserve supersymmetry, as discussed in the text.

Appendix B

This appendix contains the expression of the parameter R defined in eq. (28), which can be derived from the results of ref. [16]. The SM contribution is

$$\Delta_{SM} = \frac{tw^3 - 12tw^2 + 15tw - 4 + 6tw^2 \ln tw}{4(tw - 1)^3} \quad (\text{B1})$$

with $tw = \bar{m}_t(\mu_w)^2/M_w^2$. The supersymmetric contribution can be split into charged Higgs and chargino contributions:

$$\Delta_{SUSY} = \Delta_H + \tilde{\Delta}. \quad (\text{B2})$$

$$\Delta_H = \cot^4 \beta \frac{(th^2 - 1 - 2th \ln th) th}{4(th - 1)^3} + 2 \cot^2 \beta tw \left[F'(tw, hw) + \frac{1}{4} G'(tw, hw) \right] \quad (\text{B3})$$

with $th = \bar{m}_t(\mu_w)^2/M_{H^\pm}^2$ and $hw = M_{H^\pm}^2/M_w^2$. The functions F' and G' are given by

$$\begin{aligned} F'(x, y) &= \frac{(x^2 - y) \ln x}{(x - y)^2(x - 1)^2} - \frac{y \ln y}{(x - y)^2(y - 1)} - \frac{1}{(x - y)(x - 1)} \\ G'(x, y) &= \frac{x}{(x - y)(x - 1)} \left[1 - \left(\frac{1}{x - 1} + \frac{y}{x - y} \right) \ln x \right] + \frac{y^2 \ln y}{(x - y)^2(y - 1)} \end{aligned} \quad (\text{B4})$$

Consistently with our assumptions, we keep only the light stop contributions to $\tilde{\Delta}$:

$$\tilde{\Delta} = \sum_{i,j=1,2} \frac{M_w^4}{m_t^2 m_{\chi_j}^2} t_{2i}^2 t_{2j}^2 G'(y_{2j}, x_{ij}), \quad (\text{B5})$$

where y_{2j} is the variable defined in eq. (7) and x_{ij} is the ratio of the squared masses of two charginos.

References

- [1] M.S. Alam *et al.* (CLEO Collab.), *Phys. Rev. Lett.* **74** (1995) 2885.
- [2] S. Glenn (for the CLEO Coll.), talk at the Meeting of the American Physics Society, Columbus, Ohio, March 1998, CLEO-TALK 98-4.
- [3] R. Barate *et al.* (ALEPH Coll.), preprint CERN-EP/98-044.

- [4] K. Chetyrkin, M. Misiak and M. Münz, *Phys. Lett.* **B400** (1997) 206.
- [5] A. Ali and C. Greub, *Phys. Lett.* **B361** (1995) 146;
C. Greub, T. Hurth and D. Wyler, *Phys. Lett.* **B380** (1996) 385; *Phys. Rev.* **D54** (1996) 3350.
- [6] K. Adel and Y.P. Yao, *Phys. Rev.* **D49** (1994) 4945;
C. Greub and T. Hurth, *Phys. Rev.* **D56** (1997) 2934;
A.J. Buras, A. Kwiatkowski, N. Pott, *Nucl. Phys.* **B517** (1998) 353.
- [7] M. Ciuchini, G. Degrossi, P. Gambino and G.F. Giudice, preprint CERN-TH/97-279, MPI-PhT/97-64, hep-ph/9710335 to appear on *Nucl. Phys.* **B**.
- [8] M.B. Voloshin, *Phys. Lett.* **B397** (1997) 295;
A. Khodjamirian, R. Rückl, G. Stoll, and D. Wyler, *Phys. Lett.* **B402** (1997) 167;
Z. Ligeti, L. Randall, and M.B. Wise, *Phys. Lett.* **B402** (1997) 178;
A.K. Grant, A.G. Morgan, S. Nussinov, and R.D. Peccei, *Phys. Rev.* **D56** (1997) 3151;
G. Buchalla, G. Isidori, and S.J. Rey, *Nucl. Phys.* **B511** (1998) 594.
- [9] A. Czarnecki and W.J. Marciano, preprint BNL-HET-98/11, hep-ph/9804252.
- [10] A. Kagan and M. Neubert, preprint CERN-TH/98-99, hep-ph/9805303.
- [11] A. Ali and C. Greub, *Phys. Lett.* **B293** (1992) 226;
A.J. Buras, M. Misiak, M. Münz and S. Pokorski, *Nucl. Phys.* **B424** (1994) 374;
M. Ciuchini *et al.*, *Phys. Lett.* **B334** (1994) 137.
- [12] A.J. Buras, A. Kwiatkowski and N. Pott, *Phys. Lett.* **B414** (1997) 157.
- [13] A. Ali and C. Greub, *Phys. Lett.* **B259** (1991) 182.
- [14] P. Ciafaloni, A. Romanino, and A. Strumia, preprint hep-ph/9710312.
- [15] F. Borzumati and C. Greub, preprint ZU-TH 31/97, hep-ph/9802391.
- [16] S. Bertolini, F. Borzumati, A. Masiero, and G. Ridolfi, *Nucl. Phys.* **B353** (1991) 591.
- [17] R. Barbieri and G.F. Giudice, *Phys. Lett.* **B309** (1993) 86.

- [18] M. Misiak, S. Pokorski, and J. Rosiek, in “Heavy Flavours II”, eds. A.J. Buras and M. Lindner, World Scientific, Singapore 1998 [hep-ph/9703442]; J.L. Hewett, preprint SLAC-PUB-7774 [hep-ph/9803370].
- [19] A. Masiero and L. Silvestrini, preprint hep-ph/9711401.
- [20] M. Dine and A.E. Nelson, *Phys. Rev.* **D48** (1993) 1277; M. Dine, A.E. Nelson, and Y. Shirman, *Phys. Rev.* **D51** (1995) 1362; M. Dine, A.E. Nelson, Y. Nir, and Y. Shirman, *Phys. Rev.* **D53** (1996) 2658.
- [21] G.F. Giudice and R. Rattazzi, preprint hep-ph/9801271.
- [22] G. Altarelli, R. Barbieri, and F. Caravaglios, *Int. Jour. Mod. Phys.* **A13** (1998) 1031 [hep-ph/9712368]; J. Erler and D. Pierce, hep-ph/9801238; A. Djouadi *et al.*, *Phys. Rev.* **D57** (1998) 4179; P.H. Chankowski and S. Pokorski. *Nucl. Phys.* **B475** (1996) 3.
- [23] H. Anlauf, *Nucl. Phys.* **B430** (1994) 245.
- [24] S.P. Martin and M.T. Vaughn, *Phys. Lett.* **B318** (1993) 331.
- [25] W. Siegel, *Phys. Lett.* **B84** (1979) 193.
- [26] E. Gabrielli and G.F. Giudice, *Nucl. Phys.* **B433** (1995) 3.
- [27] A. Strumia, preprint IFUP-TH-16/98, hep-ph/9804274.
- [28] P.H. Chankowski, *Phys. Rev.* **D41** (1990) 2877; H.-C. Cheng, J.L. Feng, and N. Polonsky, *Phys. Rev.* **D56** (1997) 6875; L. Randall, E. Katz, S. Su, preprint hep-ph/9801416.




Figure 1: Dependence of the prediction for the $B \rightarrow X_s \gamma$ branching ratio on the heavy mass scale $\mu_{\tilde{g}}$ for the following choice of parameters: $\tan \beta = 1$, $M_{H^\pm} = m_{\tilde{t}_2} = m_{\chi_2} = 100$ GeV, $m_{\chi_1} = 300$ GeV, $\theta_{\tilde{t}} = -\pi/10$, $A_b = A_t$, all heavy particle masses equal to $\mu_{\tilde{g}}$; the lighter chargino is predominantly higgsino. The value corresponding to "LOdecoupled" is the result of the LO calculation in the limit in which the heavy masses decouple. The two upper lines are the LO and NLO predictions in the two-Higgs doublet model (2HDM).




Figure 2: Upper bounds on the lighter chargino and stop masses from the CLEO 95% CL limit on $\text{BR}(B \rightarrow X_s \gamma)$ in the case $M_{H^\pm} = 100$ GeV. We have taken $|\theta_t| < \pi/10$, $|\mu| < 500$ GeV, $A_b = A_t$, and set all heavy masses to 1 TeV. For $\tan\beta = 2$ and 4 we show the results of the LO and NLO calculations. The result of neglecting the new NLO supersymmetric contributions to the Wilson coefficients is labelled as "NLO running".